


PROTOCOL	INDICATION(S)	CONDITION(S)	CONTRAINDICATIONS	TREATMENT	MAX DOSES	DOSE INTERVAL
VSA	Patient VSA	30 Days of age or Grater *	Valid Ministry of health DNR	AED / S-AED	9 **	2 min
	*For patients 30 days to age 8 use Pediatric pads / attenuator pads if equipped and supplied by your Department / Service.					
	Employ every effort to maintain high quality CPR with few interruptions in compressions.					
	**If a patient is VSA and has a return of pulse & re arrests, the maximin number of doses will reset.					
OPIOID OVERDOSE	Unconscious Resp. depression or not breathing Inability to adequately ventilate Suspected opioid overdose OR Patient VSA Suspected opioid overdose	12 y/o or greater	Known Allergy to Naloxone	Administer Naloxone (Narcan) nasal spray	3	10 min
		Unconscious				
		Respiratory depression or Respiratory rate less than 10min				
ANAPHYLAXIS	Exposure to possible allergen	Anaphylaxis	None	ADULT:	2	10 min
				Age: 8y/o + Weight: 30kg + Dose: EpiPen 0.30mg		
	Signs and symptoms consistent with anaphylaxis			PEDIATRIC	2	10 min
				Age: 2y/o - 8y/o Weight: 15kg - 30kg Dose: EpiPen 0.15mg		

CATEGORY	ADULT	CHILD	INFANT	NEONATE	NOTES
AGE:	Puberty (14-16y/o)	Puberty - 1y/o	1y/o - 30 days	30 days - newborn	Compression rate is 100-120/min
LOC	Unresponsive				
Recognition	No breathing or Gaspings	No breathing / Gaspings HR less 60 BPM	No breathing / Gaspings HR less 60 BPM	No breathing / Gaspings HR less 60 BPM	Employ perishock pause on all analyzes
Assessment	Circulation-Airway-Breathing				Ventilations are 1 second in duration. 1 second in 1second out. Resume compressions immediately after 2nd breath.
Compression Rate	100 - 120 / min				Allow the chest to fully recoil
Compression Depth	2 inch or Greater	1/3 Depth of Chest (2")	1/3 Depth of Chest (1.5")	1/3 Chest (Hands Encircling Chest)	Minimize interruptions in chest compressions to less than 10 seconds
*** Minimize Interruptions in Chest Compressions ***					
Compression - Ventilation Ratio	30-2 1 & 2 Rescuer	30-2 Single Rescuer 15-2 Two Rescuer		3 to 1 1 & 2 Rescuer	Child Pads (if used by Dept.) Front & Back placement
Airway	Head Tilt Chin Lift (Modified Jaw Thrust if Suspected C-Spine injury)			Lightly tilt head back. Drop head back	
Ventilations with Advanced Airway	1 breath every 6-8 sec (8-10 breaths / min)				
	No interruptions in chest compression for ventilation				
Defibrillation	Attach AED A.S.A.P.			No S-AED defibrillation	
	Adult Pads	Child Pads (if used by Dept.) Adult Pads			